

PA 1240962

PCT/IB04/03815

REC'D 22 NOV 2004

WIPO PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 28, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.


APPLICATION NUMBER: 60/526,546

FILING DATE: December 02, 2003

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



T. Lawrence

T. LAWRENCE
Certifying Officer

17707 U.S. PTO

PTO/SB/16(06-03)

Approved for use through 7/31/2003. OMB 0651-0032

U.S. Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Expr ss Mail Lab IN .

INVENTOR(S)

Given Name (first and middle (if any))	Family Name or Surname	Residence (City and either State or Foreign Country)
Thomas A.	Brandt	North Stonington, CT, USA

Proc ss For Converting Heterocyclic Ketones To Amido-Substituted Heterocycles

Direct all correspondence to:

CORRESPONDENCE ADDRESS

☒ Customer Number

28523

☐ Firm or
Individual Name

Address

Address

City

State

Zip

Country /

Telephone

Fax

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification

Number of Pages

15

☐ CDs, Number

☐ Drawing(s)

Number of Sheets

☐ Other (specify)

☒ Application Data Sheet. See 37 CFR1.76

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

**FILING FEE
AMOUNT(\$)**

\$160.00

☐ Applicant claims small entity status. See 37 CFR1.27

☐ A check or money order is enclosed to cover the filing fees

☒ The Director is hereby authorized to charge all required
filing fees to, and credit any overpayment to Deposit Account Number:

16-1445

☐ Payment by credit card Form PTO-2038 is attached.

The invention was made by an agency of the United States Government or under a contract with an agency of the
United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE

Arlene K. Musser

DATE:

December 2, 2003

TYPED or PRINTED NAME

Arlene K. Musser

REGISTRATION NO 37,895

(if appropriate)

TELEPHONE

860-715-0871

Docket Number.

PC26214

**USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT
ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROCESS FOR CONVERTING HETEROCYCLIC KETONES
TO AMIDO-SUBSTITUTED HETEROCYCLES

5 The present invention relates to an improved process for preparing amido-substituted 4- to 6-membered heterocyclic compounds from 4- to 6-membered heterocyclic ketones. The amido-substituted 4- to 6-membered heterocyclic compounds are useful intermediates in the synthesis of cannabinoid (CB-1) antagonists.

10


BACKGROUND

 The synthesis of α -amino acids by reaction of aldehydes with ammonia and hydrogen cyanide followed by hydrolysis of the resulting α -aminonitriles is known as the Strecker Amino-Acid Synthesis. See, A.
15 Strecker, *Ann*, **75**, 27 (1850); and A. Strecker, *Ann*, **91**, 349 (1854). Over the years, safer, milder, and more selective reaction conditions have been developed, especially in regard to asymmetric synthesis. In addition, the scope of the reaction has been extended to include primary and secondary amines. See, e.g., J. P. Greenstein, M. Winitz, *Chemistry of the Amino*
20 *Acids*, vol. 3 (New York, 1961) pp 698-700; G.C. Barrett, "Asymmetric synthesis using enantiopure sulfinimines", *Chemistry and Biochemistry of the Amino Acids* (Chapman and Hall, New York, 1985) pp 251, 261.; F.A. Davis, et al., "Review of Stereoselective Synthesis", *Tetrahedron Letters*,
25 **35**, 9351 (1994); R.O. Duthaler, *Tetrahedron*, **50**, 1539-1650 *passim* (1994).

 Although the Strecker reaction provides a convenient means for making α -aminonitriles, the use of cyanide reagents raises safety issues due to the high toxicity of any residual cyanide in the reaction mixture. Therefore, there is a need for an efficient means for producing an α -
30 aminoamide from the corresponding α -aminonitrile without the risk of exposure to residual cyanide from the preparation of the intermediate α -aminonitrile.

SUMMARY

The present invention provides a process for preparing a compound of Formula (I) having little or no risk of exposure to residual cyanide.



wherein

R^{4b} and $R^{4b'}$ are each independently hydrogen or (C_1-C_6) alkyl;

X is a bond, $-CH_2CH_2-$ or $-C(R^{4c})(R^{4c'})-$, where R^{4c} and $R^{4c'}$ are each independently hydrogen or (C_1-C_6) alkyl;

R^{4d} is hydrogen, (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, or taken together with $R^{4d'}$ forms a 4- to 6-membered heterocyclic ring optionally containing an additional heteroatom selected atom N, O, or S;

$R^{4d'}$ is hydrogen, (C_1-C_6) alkyl, or taken together with R^{4d} forms a 4- to 6-membered heterocyclic ring optionally containing an additional heteroatom selected from N, O or S;


Z is a bond, $-CH_2CH_2-$, or $-C(R^{4e})(R^{4e'})-$, where R^{4e} and $R^{4e'}$ are each independently hydrogen or (C_1-C_6) alkyl; and

R^{4f} and R^{4r} are each independently hydrogen or (C_1-C_6) alkyl;

or a pharmaceutically acceptable salt thereof;


comprising the steps of

(1) reacting a compound having a formula $R^{4d}-NH-R^{4d'}$ and a cyanide source with a compound of Formula (Ia) to form an intermediate of Formula (Ib)



where Pg is an amino-protecting group and R^{4b} , $R^{4b'}$, X, Z, R^{4d} , $R^{4d'}$, R^{4f} and $R^{4f'}$ are as defined above;

- 5 (2) hydrolyzing the nitrile group of the compound of Formula (Ib) with alkaline hydrogen peroxide in the presence of dimethylsulfoxide to form a compound of Formula (Ic)



- 10 where Pg, R^{4b} , $R^{4b'}$, X, Z, R^{4d} , $R^{4d'}$, R^{4f} and $R^{4f'}$ are as defined above;

 (3) removing the amino-protecting group to form the compound of Formula (I); and

 (4) optionally forming a pharmaceutically acceptable salt of said compound of Formula (I).

- 15 Preferably, the compound of Formula (Ia) is converted to the compound of Formula (Ic) without isolating the compound of Formula (Ib). For the compounds of Formula (I) and corresponding intermediates, R^{4b} , $R^{4b'}$, R^{4f} , $R^{4f'}$ are preferably all hydrogens. X is preferably $-\text{CH}_2-$ or a bond. Z is preferably $-\text{CH}_2-$ or a bond (more preferably, X and Z are both a bond).
- 20 R^{4d} is preferably $(\text{C}_1\text{-C}_6)\text{alkyl}$ (more preferably, R^{4d} is ethyl) and $R^{4d'}$ is preferably hydrogen.

Definitions

As used herein, the term "alkyl" refers to a hydrocarbon radical of the general formula C_nH_{2n+1} . The alkane radical may be straight or branched. For example, the term "(C₁-C₆)alkyl" refers to a monovalent,
5 straight, or branched aliphatic group containing 1 to 6 carbon atoms (e.g., methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl, *t*-butyl, *n*-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, neopentyl, 3,3-dimethylpropyl, hexyl, 2-methylpentyl, and the like). Similarly, the alkyl portion (i.e., alkyl moiety) of an alkylamino group has the same definition as above. The
10 term "di(C₁-C₆)alkyl" refers to two (C₁-C₆)alkyl groups which may be the same or different.

The term "cycloalkyl" refers to a carbocyclic ring system which may include alkyl substitutions. For example, (C₃-C₆)cycloalkyl includes cyclopropyl, methylcyclopropyl, cyclobutyl, methylcyclobutyl,
15 dimethylcyclobutyl, cyclopentyl, methylcyclopentyl, and cyclohexyl.

The term "cyanide source" refers to any reagent that can provide a cyanide ion under the reaction conditions. For example, potassium cyanide, sodium cyanide, trimethylsilyl cyanide, hydrogen cyanide, and the
like.

20 The phrase "pharmaceutically acceptable" indicates that the substance or composition must be compatible chemically and/or toxicologically, with the other ingredients.

The term "protecting group" or "Pg" refers to a substituent that is commonly employed to block or protect a particular functionality while
25 reacting other functional groups on the compound. For example, an "amino-protecting group" is a substituent attached to an amino group that blocks or protects the amino functionality in the compound. A preferred amino-protecting is benzhydryl.

30 DETAILED DESCRIPTION

The process of the present invention provides a convenient and efficient means for preparing intermediates that are useful in making compounds that have been found to be cannabinoid (CB-1) antagonists.

10

The reaction scheme illustrates the synthesis of 1,4-dihydropyridine-3-carbonitriles (I) from 2-hydroxy-1,2,3,4-tetrahydropyridines. The process begins with a 2-hydroxy-1,2,3,4-tetrahydropyridine derivative (where the nitrogen is substituted with a protecting group 'Pg' and the ring contains substituents 'X' and 'Z'). This intermediate undergoes cyclization to form a 2-oxo-1,2,3,4-tetrahydropyridine derivative (labeled 1(a)). Intermediate 1(a) then reacts with a diamine (R^{4d}-N-R^{4d}) to form a 1,4-dihydropyridine-3-carbonitrile derivative (labeled 1(b)). Intermediate 1(b) is shown in brackets, indicating it is a reactive intermediate. Finally, 1(b) is converted to the final product (I), which is a 1,4-dihydropyridine-3-carbonitrile derivative where the nitrogen is substituted with a hydrogen atom (H) and the ring contains substituents 'X' and 'Z'.

15

20

procedures. For example, the hydroxy compound may be treated with oxalyl chloride and dimethyl sulfoxide in the presence of a base (e.g., triethylamine) to form the ketone 1(a) (also known as the Swern oxidation). The ketone 1(a) is reacted with the desired amino compound (R^{4d} -NH- $R^{4d'}$, where R^{4d} and $R^{4d'}$ are as defined above) and a cyanide source in a protic solvent (e.g., methanol and/or water) to form the nitrile 1(b). Suitable amino compounds include alkylamines (e.g., methyl amine, ethyl amine, *n*-propylamine, *iso*-propyl amine, *n*-butylamine, *sec*-butylamine, *iso*-butyl amine, and the like.), dialkylamines (e.g., dimethylamine, diethylamine, methylethylamine, and the like), cycloalkylamines (e.g., cyclopropylamine, methylcyclopropylamine, cyclobutylamine, methylcyclobutylamine, dimethylcyclobutylamine, cyclopentylamine, methylcyclopentylamine, cyclohexylamine, and the like), and heterocyclic amines (e.g., azetidine, pyrrolidine, imidazolidine, oxazolidine, thiazolidine, piperidine, piperazine, morpholine, thiamorpholine, and the like). When a cyanide salt is used for the cyanide source, then the reaction medium needs to be acidic for the generation of hydrogen cyanide. For example, acetic acid or hydrochloric acid is typically added with potassium cyanide. The nitrile intermediate 1(b) is then hydrolyzed to the amide 1(c) using procedures analogous to those described by Yasuhiko Sawaki and Yoshiro Ogata in Bull Chem Soc Jpn, **54**, 793-799 (1981). For example, nitrile intermediate 1(b) is treated with about 1.1 equivalents of alkaline hydrogen peroxide (e.g., hydrogen peroxide in the presence of a strong base (e.g., sodium hydroxide or potassium hydroxide) in the presence of about 1.2 equivalents of dimethylsulfoxide (DMSO) in a protic solvent (e.g., methanol). Generally, the amount of sodium hydroxide added is about 3 mol% over the amount of total acid used in the Strecker reaction (e.g., mol acetic acid plus mol HCl from amine hydrochloride salt) The pH is about 13. Preferably, the hydrolysis to the amide 1(c) is performed with the crude reaction mixture from the previous step without isolating the α -aminonitrile intermediate 1(b). Finally, the protecting group may be removed using procedures appropriate for the particular protecting group utilized. For example, when

benzhydryl is the protecting group, it may be removed by hydrogenation in the presence of a catalyst (e.g., $\text{Pd}(\text{OH})_2$).

There are several advantages of the process of the present invention over other processes that could be used for this conversion. For example, the introduction of the nitrile group into the molecule and the subsequent hydrolysis to the amide can be done in a single pot reaction. When X and Z are both a bond and R^{4d} is ethylamino, the amide 1(c) was isolated directly from the crude reaction mixture in sufficient purity to be used in the next step without any further purification, thus providing an efficiency advantage in manufacturing. In addition, the oxidizing agent (basic hydrogen peroxide) likely decomposes any remaining cyanide, presumably to cyanate and then further to carbon dioxide and ammonia, thus eliminating the safety issue associated with cyanide exposure and waste stream management. The use of basic peroxide hydrolysis allowed the reaction to take place in the presence of amine functionality which under neutral or slightly acidic H_2O_2 would likely have oxidized the tertiary amine to an N-oxide and the secondary amine to an oxime. In the present invention, the rate of nitrile hydrolysis is essentially instantaneous such that oxidative side reactions are relatively slow if present at all.

EXAMPLES

Unless specified otherwise, starting materials are generally available from commercial sources such as Aldrich Chemicals Co. (Milwaukee, WI), Lancaster Synthesis, Inc. (Windham, NH), Acros Organics (Fairlawn, NJ), Maybridge Chemical Company, Ltd. (Cornwall, England), Tyger Scientific (Princeton, NJ), and AstraZeneca Pharmaceuticals (London, England).

General Experimental Procedures

NMR spectra were recorded on a Varian Unity™ 400 or 500 (available from Varian Inc., Palo Alto, CA) at room temperature at 400 and 500 MHz ^1H , respectively. Chemical shifts are expressed in parts per million (δ) relative to residual solvent as an internal reference. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; q, quartet; m,

multiplet; br s, broad singlet; v br s, very broad singlet; br m, broad multiplet; 2s, two singlets. In some cases only representative ^1H NMR peaks are given.


Mass spectra were recorded by direct flow analysis using positive and negative atmospheric pressure chemical ionization (APCI) scan modes. A Waters/APCI/MS model ZMD mass spectrometer equipped with Gilson 215 liquid handling system was used to carry out the experiments.

Mass spectrometry analysis was also obtained by RP-HPLC gradient method for chromatographic separation. Molecular weight identification was recorded by positive and negative electrospray ionization (ESI) scan modes. A Waters/Micromass ESI/MS model ZMD or LCZ mass spectrometer equipped with Gilson 215 liquid handling system and HP 1100 DAD was used to carry out the experiments.

1-Benzhydryl-azetidin-3-ol is available from DCI Pharmtech, Inc. (Taiwan)

Example 1

Preparation of 1-Benzhydryl-azetidin-3-one (I-1a):




Oxalyl chloride (145.2 g, 1.121 mol) was added to dichloromethane (3.75 liters) and the resulting solution was cooled to -78°C . Methyl sulfoxide (179.1 g, 2.269 mol) was then added over a duration of 20 minutes (maintained internal temperature $<-70^{\circ}\text{C}$ during addition). 1-Benzhydryl-azetidin-3-ol (250.0 g, 1.045 mol) was then added as a solution in dichloromethane (1.25 liter) to -78°C solution over a duration of 40 minutes (maintained internal temperature $<-70^{\circ}\text{C}$ during addition). The solution was stirred for 1 hour at -78°C followed by the addition of triethylamine (427.1 g, 4.179 mol) over 30 minutes (maintained internal temperature $<-70^{\circ}\text{C}$ during addition). Reaction was then allowed to come

to room temperature slowly and stir for 20 hours. 1.0 M hydrochloric acid (3.2 liters, 3.2 mol) was added to the crude reaction solution over 30 minutes, followed by stirring for 10 minutes at room temperature. The heavy dichloromethane layer (clear yellow in color) was then separated and discarded. The remaining acidic aqueous phase (clear, colorless) was treated with 50% sodium hydroxide (150 ml, 2.1 mol) with stirring over a 30 minute period. The final aqueous solution had a pH=9. At this pH, the desired product precipitates from solution as a white solid. The pH=9 solution was stirred for 30 minutes and then the precipitated product was collected by filtration. The collected solid was washed with 1.0 liter of water and then air dried for 36 hr to give 1-benzhydryl-azetidin-3-one (l-1a) (184.1 g, 74%) as an off-white solid.

+ESI MS (M+1) 256.3 (M+1 of hydrated ketone); ^1H NMR (400 MHz, CD_2Cl_2) δ 7.47-7.49 (m, 4H), 7.27-7.30 (m, 4H), 7.18-7.22 (m, 2H), 4.60 (s, 1H), 3.97 (s, 4H).

Preparation of 1-Benzhydryl-3-ethylamino-azetidine-3-carboxylic acid amide (l-1c):




1-Benzhydryl-azetidin-3-one l-1a (53.43 g, 0.225 mol) was dissolved in methanol (750 ml) to give a clear pale yellow solution. Ethylamine hydrochloride (20.23 g, 0.243 mol) was added in one portion as a solid (reaction solution remains clear) followed by addition of potassium cyanide (15.38 g, 0.229 mol) in one portion as a solid (potassium cyanide not very methanol soluble – suspended as white flakes). Acetic acid (14.86 g, 0.246 mol) was added followed by stirring for 2.5 hours at room temperature to give a homogenous suspension (white crystalline solids of uniform small size). LCMS showed nearly complete consumption of azetidinone starting material and a mixture of

1-benzhydryl-3-hydroxy-azetidine-3-carbonitrile (cyanohydrin) and
 1-benzhydryl-3-ethylamino-azetidine-3-carbonitrile (Strecker product). The
 reaction mixture was then warmed to 55°C and stirred for 15 hours and
 LCMS analysis showed a ~90:10 mixture of Strecker product:cyanohydrin
 (ratio appears to be an equilibrium ratio).

The crude reaction mixture was cooled to 50°C followed by the
 addition of dimethyl sulfoxide (21.10 g, 0.269 mol) and then addition of
 aqueous 2N sodium hydroxide (251 ml, 0.502 mol) over 10 minutes
 (maintained internal temperature >45°C). Re-analysis by LCMS shows all
 of the cyanohydrin was converted back to 1-benzhydryl-azetidin-3-one
 starting material to show a ratio of Strecker product:azetidinone of ~90:10).
 Solution pH equaled 13. To the basic reaction solution at 50°C was added
 11% aqueous hydrogen peroxide (80 ml, 0.247 mol) over 5 minutes while
 maintaining the internal temperature between 50 to 65°C. The product
 began to precipitate during peroxide addition, and after complete addition,
 water was added (270 ml) to help facilitate stirring. The reaction mixture
 was held at 50°C for 30 minutes then cooled to room temperature over 1
 hour, followed by stirring at room temperature for 1 hour. The precipitated
 product was collected by filtration and rinsed with 1.0 liter of water,
 followed by briefly air-drying on the filter for 1 hour. After further drying *in*
vacuo, 1-benzhydryl-3-ethylamino-azetidine-3-carboxylic acid amide (I-1c)
 was isolated as an off-white solid (55.31 g, 79% over two steps).

+ESI MS (M+1) 310.5; ¹H NMR (400 MHz, CD₃OD) δ 7.41 (d, J =
 7.1 Hz, 4H), 7.25 (t, J = 7.5 Hz, 4H), 7.16 (t, J = 7.5 Hz, 2H), 4.49 (s, 1H),
 3.44 (d, J = 8.3 Hz, 2H), 3.11 (d, J = 8.3 Hz, 2H), 2.47 (q, J = 7.1 Hz, 2H),
 1.10 (t, J = 7.3 Hz, 3H).

Preparation of 3-Ethylamino azetidine-3-carboxylic Acid Amide,
Hydrochloride Salt (I):




To a suspension of 1-benzhydryl-3-ethylaminoazetidine-3-carboxylic acid amide (I-1c; 36.1 g, 117 mmol) in methanol (560 ml) at room temperature was added concentrated aqueous HCl (19.5 ml, 234 mmol), resulting in a clear solution. To 20% Pd(OH)₂ on carbon (3.75 g) was added methanol (85 ml), followed by the methanolic solution of I-1c. The mixture was placed on a Parr® shaker and then reduced (50 psi H₂) at room temperature for 20 hours. The reaction was then filtered through Celite® and then concentrated to low volume under reduced pressure, at which point a precipitate forms. The suspension was diluted with 500 ml of methyl t-butyl ether (MTBE), stirred for an additional hour, and the precipitate collected by vacuum filtration. The solid was washed with MTBE and then dried, *in vacuo*, to afford (I) (24.8 g, 98%) as a colorless solid.

+APCl MS (M+1) 144.1; ¹H NMR (400 MHz, CD₂Cl₂) δ 4.56 (br s, 4H), 3.00 (q, *J* = 7.2 Hz, 2H), 1.36 (t, *J* = 7.1 Hz, 3H).

CLAIMS

What is claimed is:

1. A process for preparing a compound of Formula (I)



(I)

wherein

R^{4b} and $R^{4b'}$ are each independently hydrogen or (C₁-C₆)alkyl;

X is a bond, -CH₂CH₂- or -C(R^{4c})(R^{4c'})-, where R^{4c} and R^{4c'} are each independently hydrogen or (C₁-C₆)alkyl;

R^{4d} is hydrogen, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, or taken together with $R^{4d'}$ forms a 4- to 6-membered heterocyclic ring optionally containing an additional heteroatom selected atom N, O, or S;


$R^{4d'}$ is hydrogen, (C₁-C₆)alkyl, or taken together with R^{4d} forms a 4- to 6-membered heterocyclic ring optionally containing an additional heteroatom selected from N, O or S;

Z is a bond, -CH₂CH₂-, or -C(R^{4e})(R^{4e'})-, where R^{4e} and R^{4e'} are each independently hydrogen or (C₁-C₆)alkyl; and

R^{4f} and $R^{4f'}$ are each independently hydrogen or (C₁-C₆)alkyl; or a pharmaceutically acceptable salt thereof;


comprising the steps of

(1) reacting a compound having a formula R^{4d} -NH- $R^{4d'}$ and a cyanide source with a compound of Formula (Ia) to form an intermediate of Formula (Ib)



where Pg is an amino-protecting group and R^{4b} , $R^{4b'}$, X, Z, R^{4d} , $R^{4d'}$, R^{4f} and $R^{4f'}$ are as defined above;

- 5 (2) hydrolyzing the nitrile group of the compound of Formula (Ib) with alkaline hydrogen peroxide in the presence of dimethylsulfoxide to form a compound of Formula (Ic)



- 10 where Pg, R^{4b} , $R^{4b'}$, X, Z, R^{4d} , $R^{4d'}$, R^{4f} and $R^{4f'}$ are as defined above;

(3) removing the amino-protecting group to form the compound of Formula (I); and

(4) optionally forming a pharmaceutically acceptable salt of said compound of Formula (I).

15

2. The process of Claim 1 wherein said compound of Formula (Ia) is converted to said compound of Formula (Ic) without isolating said compound of Formula (Ib).

20

3. The process of Claim 2 wherein R^{4b} , $R^{4b'}$, R^{4f} , $R^{4f'}$ are all hydrogens.

4. The process of Claim 3 wherein X is -CH₂- or a bond; and Z is -CH₂- or a bond.


5 5. The process of Claim 4 wherein R^{4d} is (C₁-C₆)alkyl and R^{4d'} is hydrogen.

6. The process of Claim 5 wherein X and Z are both a bond.

10 7. The process of Claim 5 or 6 wherein R^{4d} is ethyl.

ABSTRACT

The present invention provides a safe and convenient process for preparing compounds of Formula (I) from the corresponding heterocyclic ketone.



(I)